

Neutron spin-dependent structure function, Bjorken sum rule, and first evidence for singlet contribution at low x

J. Soffer¹, and O. Teryaev²

(1) *Centre de Physique Théorique - CNRS - Luminy,
Case 907 F-13288 Marseille Cedex 9 - France*

(2) *Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, Dubna, 141980, Russia*

ABSTRACT

We perform the isospin decomposition of proton and neutron SLAC data in the region $0.01 \leq x \leq 0.1$. The isovector part is well described by a power behaviour x^α , where α leads to the validity of Bjorken sum rule and it is consistent with the power extracted from all previous data using NLO evolution. The isoscalar part behaviour may be interpreted as a partial cancellation between a positive non-singlet contribution and a singlet one strongly negative.

The very accurate measurement of the neutron spin-dependent structure function $g_1^n[1]$, whose results have been presented recently, possess two remarkable properties. First, the values of neutron structure function at $Q^2 = 5\text{GeV}^2$ are rather large and negative in the region of moderately low x . Second, the data can be rather accurately fitted by the power function $x^{-0.8}$ and this power seems to be unexpectedly large. It is not obvious, where this large number, for which there is no indication in the proton data, is coming from, so it may affect the extrapolation to $x = 0$ and cast some doubts on the validity of Bjorken sum rule. In the present work we perform the isospin decomposition of the data for both proton and neutron. As a result, we conclude that the isovector contribution is well approximated by the power behaviour found earlier by an elaborate method based on next-to-leading order (NLO) evolution[2]. It may be also interpreted as the manifestation of $\ln^2 x$ terms[3]. For the isoscalar part one has the signature for a rather singular singlet contribution, compatible with the similar behaviour predicted by QCD[4]. This may imply a significant gluon polarization in the nucleon in this range of x .

The main feature of the new neutron data[1] is large and negative $g_1^n \sim -g_1^p$, measured with a good accuracy up to small x , say $x \sim 0.01$. The SLAC proton data[5] at $Q^2 = 3\text{GeV}^2$ are positive and of roughly the same magnitude, but on the contrary, rather flat in this region. This could be understood, qualitatively, as a result of the interplay of a negative contribution at low x , responsible for the singular behaviour of the neutron, and a positive contribution at larger x . To check this assumption it is instructive to consider the isovector contribution. Since there is no clear evidence of scaling violations between $Q^2 = 3\text{GeV}^2$ and $Q^2 = 5\text{GeV}^2$ in any polarized deep inelastic scattering experiment, we may neglect, for the time being, the effects of QCD evolution, because we are only interested in the results,

provided by the data at the present level of statistical accuracy. It is then possible, by combining the SLAC neutron data with the SLAC proton data, to determine the quantity $g_1^{p-n} \equiv g_1^p - g_1^n$, entering the Bjorken sum rule, with a higher accuracy, than for proton alone. This is because the neutron data are fortunately negative, so the difference is larger in magnitude than g_1^p , while the errors are practically the same for g_1^{p-n} and g_1^p , given the quality of the neutron data.

Performing this analysis of the data, we see immediately a behaviour less sharp than that of neutron. Since phase space effects, as powers of $(1-x)$, are not so important in the region under consideration, we were looking for a simple power parametrization of the data, implied by Regge pole behaviour, and we found that this works rather well, for $0.016 \leq x \leq 0.125$ with the following expression

$$g_1^{p-n}(x) = 0.147x^{-0.45}, \quad (1)$$

as shown in Fig.1.

The power we obtain is significantly smaller than the expected contribution of the $a_1(1260)$ meson trajectory (~ 0.14). As a result, the contribution to the Bjorken integral from the region $0 \leq x \leq 0.125$ is large

$$\int_0^{0.125} dx g_1^{p-n}(x) = 0.085. \quad (2)$$

The region of higher x corresponds to a neutron contribution much smaller than that of the proton, the latter also providing a large contribution to the above integral, which is 0.09 for $0.125 < x < 1$. The total contribution to the Bjorken sum rule, obtained in this way, i.e. 0.175, appears to be in the fair agreement with the theoretical value. Note that when the final neutron data will be available, as well as more precise proton data, it will allow a more serious analysis, taking also into account the effects of QCD evolution.

At the present moment we would like to stress, that by combining the current neutron and proton data, one is led to good agreement with Bjorken sum rule. Let us stress that this power is compatible with the one obtained using NLO fit[2] to all previous data (-0.56 ± 0.21). It is consistent with the result of $\ln^2 x$ summation[3].




Figure 1: Comparison for g_1^{p-n} between the curve given by eq.(1) and the SLAC data refs.[1,5].

Since we found, that the sharp neutron structure function is not seen in the difference between proton and neutron, it should be attributed to the isoscalar channel. Also, the partial cancellation, we suspect to be at the origin of a flat proton structure function, should be manifested in this channel as well. To check this, we calculated the quantity $g_1^{p+n} \equiv g_1^p + g_1^n$. It really shows a rather flat structure for $x \geq 0.035$ which should be related to the interplay of the negative sharp contribution, showing itself in the fit $x^{-0.8}$ to the neutron data[1], and a positive contribution with a smaller power, dominating at larger x . Since there is no counterpart for such a sharp behaviour in the conventional Regge analysis, we make a strong, but natural assumption, that it manifested itself in the $SU(3)$ -singlet channel. It is in this channel that a strong mixing between polarized quarks and gluons provides the anomalous gluon contribution to the first moment of g_1 [6].

At low x the quark-gluon mixing provides a strong correction to the subleading behaviour[4], producing a power close to 1. For a first estimate we find that the data are well described by

$$g_1^{p+n}(x) = 0.145x^{-0.45} - 0.03x^{-0.87}, \quad (3)$$

as seen in Fig.2.

This formula is suggesting that the isoscalar contribution is approximately equal to the isovector one, which is not so surprising, in order to have the neutron structure function, dominated by the most singular power only. Note that eq.(3) gives also a fair description of the g_1^d data in this kinematical region[7] and further details on this result are given in ref.[8].




Figure 2: Same as Fig.1 for g_1^{p+n} with the curve given by eq.(3).

- [1] E. Hughes, (SLAC experiment E154), presented at ICHEP Warsaw, July 1996.
- [2] R.D. Ball, S. Forte and G. Ridolfi, Phys. Lett. **B378** (1996) 255; hep-ph/9608399.
- [3] J. Bartels, B.I. Ermolaev and M.G. Ryskin, Z. Phys. **C70** (1996) 273.
- [4] J. Bartels, B.I. Ermolaev and M.G. Ryskin, hep-ph/9603204, March 1996.
- [5] K. Abe et al. (SLAC experiment E143), Phys. Rev. Lett. **74** (1995) 346.
- [6] A.V. Efremov, J. Soffer, O.V. Teryaev, Nucl. Phys. **B346** (1990) 97.
- [7] K. Abe et al. (SLAC experiment E143), Phys. Rev. Lett. **75** (1995) 248.
- [8] J. Soffer, O.V. Teryaev, Preprint CPT-96/P.3387, hep-ph/9609329.